

identification of vulnerable Communities with High Lead Exposures Residence in Michigan

Hanging Cur., Valente Zartarian, Antonios Poulskos, Lindsoy Stanet

Heliona Exposes: Bessect Libertess (Blood Seventh and Development U.S. Embrumental Protection Agency Research Thangs Park, RC-ASRC Federal Ostropic Companies Eastern MA

Background and Objectives

Lead exposure is associated with numerous harmful effects in children and adults, including adverse neurological, cardiovascular, kidney, developmental, and reproductive outcomes. Exposure to even small amounts of lead can be harmful to a child's developing brain, resulting in detrimental effects on IQ and behavior. Over the past 50 years, lead exposure for the U.S. population has decreased, resulting in lowered blood lead levels. However, there are still likely to be vulnerable communities (i.e., those overburdened with few resources and disproportionately impacted by lead exposures) where significant sources of lead remain that need to be identified to prevent adverse public health outcomes. As such, our objectives were to identify high risk areas in Michigan with elevated blood lead levels, determine the main contributing factors, and develop approaches for future studies to inform exposure reduction efforts.

We obtained a blood lead dataset from the state of Michigan comprised of measurements from young children (average age ~7 years old) collected over 10 years, totaling ~2.5 million data points. Each individual blood lead measurement was geocoded to determine if neighborhood clusters existed with high lead exposures. We used ARCMAP 10.3.1 ([Getis-Ord Gi*] and Gluster and outlier analysis (Anselin Local Morans II) to identify communities with high blood lead levels (hot spots) by weighting the number of people with high blood lead levels (e.g., 3.5 or 5 µg/dl) by population or total number of people for blood lead screening. General linear model (GLM) was applied to identify main contributing factors

Conclusions

- ·Rate of high-BLL decreasing in MI but slower in higher risk locations than in areas with low-BLL
- · Consistent findings for high Pb risk areas by Reuters analysis and EPA/ORD, but ORD approach provides better geographic and temporal resolution
- -- Different GIS methods yield similar results
- · Many high Pb risk areas identified in MI with higher rates than Flint
- --e.g., Detroit area
- –Only 3 census tracts in top 100 (All three in 90th to 100th percentiles) in Flint, MI
- · Housing age, minority status, income level, and education level are found to be major contributors by GLM

As an orderigeneous

The Bureau of Epidemiology and Population Health/Division of Environmental Health, Michigan Department of Health and Human Services

EPA Office of Research and Development scientists and collaborators

Disculine

The views expressed in this presentation are that of the presenters and do not necessarily represent the views and/or policies of the U.S. Environmental Protection Agency.



If S. Environmental Protection Agency Office of Resource and Development

Reuters Analyses Nationwide Childhood St.L. American Maps by Jip Code endire Centrus Tred Underlying BLL data provided by the 4.00 feather was appearing. For reuse on president rate

